

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>A61K 48/00</b>		<b>A1</b>	(11) International Publication Number: <b>WO 96/27393</b> (43) International Publication Date: 12 September 1996 (12.09.96)
(21) International Application Number: <b>PCT/US96/02681</b> (22) International Filing Date: <b>7 March 1996 (07.03.96)</b> (30) Priority Data: <b>08/400,089 7 March 1995 (07.03.95) US</b> (71) Applicant: <b>UNIVERSITY OF PITTSBURGH [US/US]; 911 William Pitt Union, Pittsburgh, PA 15260 (US).</b> (72) Inventors: <b>HUANG, Leaf; 2468 Matterhorn Drive, Wexford, PA 15091 (US). SORGI, Frank, L.; 2640 Middletown Road, Pittsburgh, PA 15205 (US).</b> (74) Agents: <b>KOKULIS, Paul, N. et al.; Cushman, Darby &amp; Cushman L.L.P., 1100 New York Avenue, N.W., Washington, DC 20005 (US).</b>			(81) Designated States: <b>AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</b>  <b>Published</b> <i>With international search report.</i>
(54) Title: <b>A DRY POWDER FORMULATION FOR GENE THERAPY</b>			
(57) Abstract <p>Gene therapy using a non-viral vector often requires the administration of large amounts of DNA or RNA dissolved in a solution of relatively small volume. The invention provides a dry powder formulation which contains a lyophilized DNA/liposome complex, a lyophilized RNA/liposome complex, a lyophilized oligonucleotide/liposome complex or a lyophilized protein/liposome complex. The dry powder is suitable for airway delivery or topical administration as an aerosol. The dry powder can be easily reconstituted with water and is active in gene transfer <i>in vitro</i> and <i>in vivo</i>. The potency of gene transfer of the dry powder was at least 50-fold higher than that of a liquid formulation of similar composition. The composition, method of preparation and method of use are described.</p>			

Quadrant EXHIBIT 2075  
Quadrant v. Nektar  
Interference 105,219

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

## A DRY POWDER FORMULATION FOR GENE THERAPY

BACKGROUND OF THE INVENTION

Gene therapy is rapidly becoming a new therapeutic modality in experimental and clinical medicine. It is desirable that a pharmaceutical formulation, containing DNA or RNA with the transgene sequence, be prepared in large quantity with pharmaceutical quality. The currently used non-viral vectors for gene therapy are mostly two-vial formulations which are mixed at the bedside prior to administration. Such an admixture formulation is acceptable but not ideal. Furthermore, the two-vial formulations are usually dilute in their concentrations of DNA, RNA, oligonucleotide or protein, necessitating the administration of solutions of relatively large volume. A highly concentrated, stable, one-vial formulation is thus more desirable than the current formulations. We describe here the composition and the method of preparation of a dry powder formulation suitable for topical and airway administration for gene therapy and other purposes. When reconstituted in an aqueous medium, it can also be administered in liquid form.

1. Field of the Invention

The present invention relates to methods and compositions suitable for gene transfer or

delivery of gene therapy. More particularly, it relates to pharmaceutical compositions in which DNA, RNA, oligonucleotides and proteins can be complexed with lipid and delivered in a concentrated and convenient form, either as a lyophilized powder or in liquid form after reconstitution in a suitable aqueous medium.

## 2. Background Information

The field of gene therapy/gene transfer is currently advancing along a number of fronts, both experimentally and clinically. Genetic material has been transferred using viral vectors, for example adenoviruses and retroviruses, as well as nonviral methods such as microinjection, calcium phosphate coprecipitation, and transfer via liposomes. (For a comprehensive review, see Morgan and Anderson, 1993.) A number of diseases and disorders are potential candidates for some type of gene therapy, if appropriate techniques can be found to deliver genetic material to target tissues and cells. In a few cases, (e.g. Cystic Fibrosis, Familial Hypercholesterolemia and Severe Combined Immunodeficiency) gene therapy has advanced to the point of clinical trials.

Cationic liposomes have been shown to be an efficient means of transfecting a wide variety of mammalian cells. Their advantages include simplicity of preparation, ability to complex a very high percentage of DNA, versatility of use with virtually any type of naked DNA or RNA (with no theoretical size limitation), the ability to transfect many different cell types, superior

transfection efficiency, and lack of immunogenicity or biohazardous activity.

Cationic liposomes composed of a mixture of DC-Chol and DOPE have been used to transfect the  
5 airway epithelial cells with CFTR cDNA in CF knock-out mice (Hyde et al., 1993; Alton et al., 1993). The same formulation has also been used in two phase I/II clinical trials for melanoma (Nabel et al., 1992) and for CF (Caplen et al., 1994).  
10 This formulation is generally regarded as effective and non-toxic and will become commercially available in the near future.

To date, however, DNA/liposome and RNA/liposome complexes have been administered in  
15 liquid form and consequently suffer from several disadvantages. Many are two-vial formulations that must be mixed shortly before administration, resulting in inconvenience in manufacture, shipping, storage and administration. In addition, it is  
20 difficult to obtain highly concentrated solutions of DNA and RNA using this technique. This may result in a need to administer large volumes of solution to achieve an optimal therapeutic effect.

#### SUMMARY OF THE INVENTION

25 This invention provides a dry powder formulation which contains a lyophilized DNA/liposome complex, suitable for gene transfer or gene therapy. Lyophilized RNA/liposome, oligonucleotide/liposome and protein/liposome  
30 complexes are also provided. The dry powder can be easily reconstituted with water and is active in gene transfer *in vitro* and *in vivo*. It is suitable

for airway delivery as an aerosol or for topical administration as a spray. The potency of gene transfer of the dry powder was at least 50-fold higher than that of a liquid formulation of similar composition. In addition, the powder can be reconstituted for delivery in liquid form.

This invention further provides a method of preparing DNA, RNA, oligonucleotides or protein in powder form suitable for therapeutic use or gene transfer.

This invention also provides an effective and efficient method of delivering DNA, RNA, oligonucleotides or protein to tissues or cells.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. A dry powder formulation of the DNA/liposome complex. 500  $\mu$ g pUCCMVCAT DNA and 500 nmol (300  $\mu$ g) of DC-Chol liposomes were mixed and allowed to complex. Lactose was added to a final concentration of 0.25 M. This solution was placed in a -80°C freezer for several hours, further frozen in liquid nitrogen, and freeze-dried for 24 hours. The resulting powder was trituated to reduce the particle size.

Figure 2. Gene transfer activity and toxicity of DNA/liposome complex reconstituted from a dry powder formulation. 2 mg pRSV-LUC DNA and 2  $\mu$ mol (1.2 mg) of DC-Chol liposomes were mixed and allowed to complex. Dextrose was added to make a 0.25 M solution with a volume of 4 ml. This mixture was freeze-dried and reconstituted with 1 ml of distilled water. This reconstituted complex was

added to CHO cells and allowed to transfect for 44 hours. Luciferase activity ( $\square$ ) and protein content ( $\diamond$ ) of the treated cells were assayed. The reconstituted DNA/liposome complex was active in gene transfer at concentrations which showed minimal toxicity, as indicated by the high protein content of the treated cells.

Figure 3. CAT activity in the treated mice. Balb/C mice weighing 20 grams were anesthetized with i.m. injections of ketamine. The dry powder formulation of the DNA-liposome complex (as shown in FIG. 1) was delivered via intra-tracheal administration. The mice were sacrificed after 40 hours and their lungs and trachea were assayed for CAT activity. Each number (1a, 1b) represents a single mouse. Lung samples (for mice 1-8) are noted as "a", while trachea samples are noted as "b". Mice 1 and 2 were instilled with 500  $\mu$ g of DNA as a fluid reconstituted from dry powder. Mice 3-7 were insufflated with maximally 10 $\mu$ g of DNA as a dry powder. Mouse 8 was an untreated control. Lanes 9 and 10 are a positive control (1 unit CAT) and a negative control (no enzyme) for the CAT reaction, respectively.

#### DETAILED DESCRIPTION OF THE INVENTION

##### MATERIALS AND METHODS

The following abbreviations are used in the specification:

DC-Chol: 3 $\beta$ [N-(N',N'-dimethylaminoethane)-carbamoyl] cholesterol

DOPE: dioleoylphosphatidylethanolamine  
CAT: chloramphenicol acetyl transferase  
CF: cystic fibrosis  
CFTR: cystic fibrosis transmembrane conductance  
5 regulator protein  
CHO cells: Chinese Hamster ovary cells

#### Plasmid DNA

Plasmid DNA was prepared by a CsCl/EtBr  
gradient method (Sambrook et al., 1989). It was run  
10 on an agarose gel to test for purity and quantitated  
by using the  $A_{260}/A_{280}$  ratio. Plasmid pUCCMVCAT was  
constructed in this lab by Dr. Hassan Farhood (see  
Farhood et al., 1995). It contains the *E. coli*  
chloramphenicol acetyl transferase (CAT) gene driven  
15 by the human cytomegalovirus early promoter and  
cloned into a pUC18 background. Plasmid pRSV-LUC  
contains a luciferase gene driven by the Rouse  
Sarcoma Virus promoter (De Wet et al., 1987).

#### Liposomes

20 DC-Chol was synthesized by the method of  
Gao and Huang (1991). DOPE was purchased from  
Avanti Polar Lipids, Inc. DC-Chol and DOPE were  
mixed in a 3:2 molar ratio, evaporated in a stream  
of nitrogen to form a thin film, and vacuum  
25 desiccated to remove any residual chloroform. The  
lipid film was rehydrated in distilled deionized  
water which had been autoclaved. The liposomes were  
allowed to hydrate overnight. DC-Chol liposomes were  
prepared by microfluidization by using an M-110S  
30 microfluidizer (Microfluidics Corp.) to an average  
diameter below 200 nm. The liposomes were filtered



through a 0.2  $\mu$ m filter to provide sterilization. The liposomes were diluted to a final concentration of 2  $\mu$ mol/ml ( 1.2 mg/ml of total lipid). Other lipids known in the art will also be effective for these purposes, with cationic lipids being particularly suitable for complexing with DNA, RNA and other negatively charged compounds.

Examples in addition to DC-Chol include DOTMA, DOSPA, DMRIE, DOTAP, DOGS, and DDAB, which may be used alone or in combination with DOPE or other compounds.

#### Preparation of DNA/Liposome Complex and Dry Powder

500  $\mu$ g of DNA and 500 nmol (300  $\mu$ g) of DC-Chol/DOPE liposomes were mixed and allowed to complex. By complex we mean an association of molecules by noncovalent bonds, such as electrostatic, ionic, hydrophobic, hydrogen bonding and others. Complexation is generally complete in a relatively short time (95% within 10 minutes using the method described here), but will occur at a slower rate in more dilute solutions. This parameter can be determined by routine experimentation by one of ordinary skill in the art, e.g. by changes in light scattering or other methods.

Lactose (or another sugar) was added to a final concentration of 0.25 M. This solution was placed in a -80°C freezer for several hours, further frozen in liquid nitrogen, and allowed to freeze-dry for 24 hours. The resulting powder was triturated to reduce the particle size. The terms "oligo-nucleotide/lipid complex", "DNA/lipid complex",

"RNA/lipid complex", and "protein/lipid complex" refer to the respective compounds that have undergone the steps of complexing and freeze-drying, with or without the addition of a cryoprotective agent. A DNA/lipid complex, for example is a lyophilized DNA/liposome complex.

As will be evident, use of the present method is contemplated for DNA, RNA (both sense and antisense), oligonucleotides, proteins and similar compounds which have all been shown to be deliverable by cationic liposomes (see, for example, Felger and Ringold, 1989 (for DNA); Malone et al., 1989 (for RNA); Debs et al., 1990 (for protein); Bennett et al., 1992, (for oligonucleotides). Any cryoprotective agent which can be freeze-dried into powder form (e.g. amino acids, urea) could also be used for the preparation of the powder.

#### Measurement of Particle Size

Freeze-dried DNA:DC-Chol liposomes were reconstituted to their original volume by addition of deionized water and shaking. The particle size was measured by using a Coulter N4 SD sub-micron particle analyzer, using a uni-modal analysis. The sample was counted for 200 seconds. The particle size of the DNA:DC-Chol liposome complex was between 150-200 nm, both prior to and after the freeze-drying process.

#### EXAMPLE 1

##### Transfection of Mammalian Cells In Vitro

2 mg of pRSV-LUC and 2  $\mu$ mol (1.2 mg) of DC-Chol liposomes were mixed and allowed to complex.

Dextrose was added to make a 0.25 M solution in a volume of 4 ml. This mixture was freeze-dried and reconstituted with 1 ml of distilled water.

The transfection was carried out in a 48 well plate in 0.5 ml as follows. An appropriate amount of the reconstituted complex was diluted into 0.5 ml of serum-free media (composed of 0.25 ml of Hanks Balanced Salt Solution and 0.25 ml of CHO-S-SFM media, both obtained from Gibco BRL), added to CHO cells (obtained from the American Type Culture Collection) and allowed to remain on the cells for 6-7 hours. The serum-free medium was then removed and replaced with 0.5 ml per well of F-12 media (also from Gibco BRL) supplemented with 10% fetal bovine serum (from Hyclone Labs), and the incubation was continued to a total time of 44 hours. For protein measurement, cells were harvested and extracted for protein which was quantitated by using a Coomassie Plus Protein Reagent (Pierce) assay.

#### 20 Luciferase Assay

CHO cells were allowed to transfect for a total time of 44 hours. The media was aspirated and the cells were washed once with a 0.9% sodium chloride solution. The cells were lysed with 100  $\mu$ l of Lysis Buffer (Promega) per well. The cell lysate was collected, briefly centrifuged, and 4  $\mu$ l was used for the luciferase assay. Luciferase Assay Buffer (Promega) was thawed and 10 ml was used to reconstitute the Luciferase Assay Substrate (Promega). The samples were loaded into an AutoLumat LB953 (Berthold) luminometer, 100  $\mu$ l of the luciferase substrate was added to the cell

lysate, and the relative light units of each sample were counted for 20 seconds. The raw data from each sample was normalized for protein content.

#### EXAMPLE 2

##### 5 Airway Delivery of Dry Powder and Reconstituted DNA/Liposome Complex

Balb/C mice weighing 20 grams were anesthetized with i.m. injections of ketamine. A small incision was made into the trachea and a 1 ml  
10 Tuberculin syringe with a 21 gauge needle was inserted. The dry powder formulation of DNA:DC-Chol liposome complex was loaded into the syringe. A 1 ml pipet tip with a rubber bulb attached to the top of the pipet tip was attached to the top of the  
15 syringe and used to blow the powder into the trachea and lungs. This procedure was repeated several times and a rise in the chest of the mouse indicated a successful insufflation.

In the case of the reconstituted  
20 DNA/liposome complex, a small volume (75-100  $\mu$ l) of distilled water was used to reconstitute 500  $\mu$ g of the dry powder formulation. A small incision was made in the neck of the mouse to expose the trachea. The entire volume (encompassing 500  $\mu$ g of the  
25 complex) was instilled over a period of one minute.

##### CAT Assay

The mice were sacrificed 40 hours after insufflation. The lungs and trachea from each mouse were removed, placed in an extraction buffer and  
30 homogenized. The sample was briefly centrifuged and

a sample of the supernatant was assayed for protein content.

The CAT assay using [<sup>14</sup>C] chloramphenicol as a substrate was done according to a modified method of Gorman et al. (1982). A sample of the supernatant was mixed with the assay reaction mixture and incubated for several hours. Ethyl acetate (1 ml) was added to the reaction mixture, vortexed briefly, and 800  $\mu$ l of the upper layer was taken and dried in a Speedvac for 1 hour. The sample was resuspended with 30  $\mu$ l of ethyl acetate and then spotted onto a TLC plate. The plate was developed in a chloroform: methanol (95:5) solution and quantified by a phosphorimager.

15 Effect of Cryoprotectant on the Particle Size and Activity of DNA/Liposome Complex

It is important in some instances, particularly for storage purposes, to include in the formulation of the invention a sufficient amount of a cryoprotectant to permit the formulation to be freeze-dried without loss of activity. Four different sugars (dextrose, sucrose, lactose, and mannitol) were tested for their potential to serve as cryoprotectants. Their ability to maintain the original size of the liposomes and the appearance of the powder mass were used as criteria to select the most appropriate sugar. As shown in Table 1, the original size of the liposomes was 177 nm. Lactose, dextrose, and sucrose were able to maintain the original size of the liposomes, while mannitol was not. Lactose produced a fine white powder upon freeze-drying, while dextrose and sucrose produced a

12

crystalline substance. Lactose was chosen as the model sugar for the dry powder formulation, although it will be appreciated that other sugars could be usefully employed.

Table 1: Size of Reconstituted Liposomes<sup>a</sup>

Cryoprotectant	Mean Diameter (nm) <sup>b</sup>				
	Cryoprotectant Concentration (M)				
	0	0.1	0.25	0.5	1.0
Lactose	1450	252	158	128	
Sucrose		171	205	181	156
Dextrose		168	273	180	301
Mannitol		765	875	985	397

<sup>a</sup> Liposomes consist of a 3:2 molar ratio of DC-Chol and DOPE in a concentration of 2  $\mu\text{mol/ml}$ . They were prepared in water and microfluidized to produce liposomes. The mean diameter of freshly prepared liposomes was 177 nm.

<sup>b</sup> The mean diameter was measured by using a Coulter N4SD submicron particle size analyzer using a unimodal analysis. The liposomes were in a volume of 2 ml prior to lyophilization and were subsequently reconstituted with water back to the original volume of 2 ml.

5           The amount of cryoprotection used in the  
present formulation can be varied depending on  
other factors. A suitable amount of the sugar or  
other cryoprotectant should usually be used to  
10 protect the DNA and liposomes during  
lyophilization. The amount is generally  
sufficient if the average size of the vesicles is  
increased or decreased by less than about 50%, and  
ideally by less than about 20%, as measured by the  
described method. A cryoprotective substance is  
15 considered to be particularly suited to these  
purposes if, in addition, it produces a powder  
with bulk and flow properties such that it can be  
administered as a dry aerosol. Identification of  
such compounds and suitable amounts or  
20 concentrations can be done by routine  
experimentation by those of ordinary skill in the  
art.

#### A Dry Powder Formulation

          A dry powder formulation prepared by  
25 lyophilizing a 1 ml solution containing 500 µg of  
DNA, 300 µg of DC-Chol liposomes and 0.25 M  
lactose is shown in FIG. 1. The powder was fine  
and flowed very well. It was stored in a  
desiccator at room temperature until use. Upon  
30 reconstitution by adding water, the DNA/liposome  
complex continued to be 150-200 nm in mean  
diameter and there was no flocculates or  
precipitates in the reconstituted solution. The  
transfection activity of the reconstituted complex  
35 was assayed with CHO cells by using complexes  
containing pRSVLUC DNA. The treated cells were  
assayed for luciferase activity two days after the



transfection. As shown in FIG. 2, cells could be effectively transfected to express the luciferase activity by the reconstituted complex in a dose dependent manner. Lack of cytotoxicity of the complex was evident because the treated cells contained nearly the same amount of protein as the untreated control (FIG. 2). Thus, we conclude the dry powder formulation, after reconstitution with water, is stable and active in gene transfer for the mammalian cells.

#### Airway Delivery of the Dry Powder

Dry powder containing pUCCMVCAT plasmid DNA and DC-Chol liposomes was insufflated intra-tracheally into mice. Up to 10  $\mu$ g of DNA was delivered into each animal. 100  $\mu$ l of reconstituted complex (500  $\mu$ g of DNA) from the same dry powder was also intra-tracheally injected as a comparison. The mice were sacrificed two days later and the major organs were removed. CAT activities of protein extracts from trachea and lung tissues of the mice were assayed by using a TLC method. As can be seen in FIG. 3, no CAT activity could be found in either the trachea or lung of the control mouse which had not been injected with any complex (lane 8). Mice treated with dry powders (lanes 3-7) had expressed CAT activity in both trachea and lung tissues. Mice injected with the reconstituted complex also expressed CAT activity in both tissues (lanes 1 and 2). It is important to note that mice treated with the dry powder received only 10  $\mu$ g of DNA maximally, yet the level of transgene expression

was similar to those treated with 500  $\mu$ g of DNA as a reconstituted fluid from the same dry powder (FIG. 3). This remarkable and surprising result indicated that the dry powder formulation was at least 50-fold more potent in activity than the reconstituted fluid.

### EXAMPLE 3

#### Treatment of Cystic Fibrosis

DNA which encodes human CFTR (prepared, e.g., as described by Caplen et al., 1994 or Caplen et al., 1995) will be complexed to DC-Chol liposomes, combined with lactose, and lyophilized, as described. The resulting powder will be delivered to the airway of a CF patient using a conventional type of dry powder inhaler (see, e.g., Atkins et al., 1992, Byron, 1990 and Hickey, 1992). In addition to CF, patients with lung cancer and other pulmonary diseases could also be treated using this method with other forms of DNA, RNA, oligonucleotides or proteins. The DNA could contain a tumor suppressor gene for cancer treatment or an antiinflammatory gene for asthma, for example. Since the dry powder is stable it lends itself to being produced in a conventional dosage form. This dosage form is preferred because it would allow patients to self-medicate as each actuation of the inhaler could deliver a predetermined dose. A patient could be treated at home as needed, eliminating the need for hospitalization or medical visits.

DISCUSSION

Since the pioneering work of Felgner et al. (1987), a number of different cationic liposome/micelle formulations have been developed, some of which are commercially available. Our work in this area started with a study of structure-activity relationships in which about 20 different cholesterol derivatives were synthesized and screened for activity and toxicity in a 1:1 mixture with dioleoyl phosphatidylethanolamine (DOPE). A set of empirical rules on the structure of cationic cholesterol was derived for the best transfection activity, best shelf stability, and least cytotoxicity (Farhood et al., 1992). DC-Chol was synthesized according to these rules and determined to be one of the best performing cationic lipids for transfection (Gao and Huang, 1991). It is known that the best transfection activity of cationic liposomes in a serum-free medium occurs at the ratio of positive/negative charge slightly greater than 1 (Gao and Huang, 1991). This is because the complex (with slightly excess positive charge) can readily bind to the negatively charged cell surface. However, most of the optimal DNA/liposome complexes lose a substantial amount of activity when serum proteins are present in the transfection medium. This is the primary reason why the cationic liposomes do not work well if injected i.v., despite some isolated reports of success (Zhu et al., 1993; Brigham et al., 1991). The ratio of DNA to liposome for airway administration varies a great deal among different investigators and ranges from

8:1 (Logan et al., 1995) to 1:8 (Alton et al., 1993) by weight. The role of proteins and other negatively charged molecules in the lining fluid of airway epithelium must play a crucial role in determining the optimal ratio of DNA/liposome for transfection. In the case of solid tumor, we found that 1:0.6 (w/w) ratio of DNA/liposome was optimal if the complex is directly injected into the tumor for transfection (unpublished data). No significant aggregation or precipitation of the DNA/liposome complex occurs at this ratio even if the concentration is greater than 1 mg of DNA per ml. A lyophilized dry powder of the complex could be prepared by using 0.25 M lactose as a cryoprotectant (see FIG. 1). Upon reconstitution, there is no change in particle size (150-200 nm in diameter) and the transfection activity is restored (FIG. 2). In vivo gene transfer activity of this powder formulation was shown by intra-tracheal administration into mice. The activity was much greater than that of the reconstituted fluid from the same dry powder.

The dry powder will preferably be administered as an aerosol in a metered dose inhaler. The powder will be suspended in Freon or another suitable agent and packaged into a sealed vial mounted onto an inhaler. Other agents that might be suitable (for example, dichlorotetrafluoroethane and dichlorodifluoromethane) are known in the art. A unit volume of the Freon suspension will be delivered to the airway by inhalation when the inhaler is triggered to dispense the suspension. Alternatively, the

powder can be loaded into a capsule which is then punctured by an inhaler to release the powder and blown into the airway. Such devices are well known in the medical arts (see, for example, 5 *Pharmaceutical Inhalation Aerosol Technology*, J. Swarbrick, ed., Marcel Decker, New York, 1992, particularly at pp. 155-185 and 255-288; and *Respiratory Drug Delivery*, P.R. Byron, ed., CRC Press, Inc., Boca Raton, Florida, 1990, 10 particularly at pp. 167-206; the entire contents of each are incorporated herein by reference). The dry powder can also be administered topically to skin or mucosal tissues by a similar delivery device.

15           The DNA/liposome complex can also be reconstituted from the dry powder and used as a liquid in cases where that would be a preferred route of administration. As the powder provides a stable product until reconstitution, any current 20 route of drug administration can be used (oral, intravenous, intramuscular, intraperitoneal, subcutaneous, topical, aerosol, etc.)

          While the invention has been described most fully in terms of formulations and uses for 25 gene transfer or therapy, it is not intended to be limited to those uses. It will be clear to the skilled artisan that formulations can also be made from a variety of suitable proteins that might be advantageous to administer diagnostically, 30 therapeutically or experimentally in the manner described. Advantages would include increased stability of the protein/lipid powder over other available forms of the protein, increased efficacy

of delivering either systemic or targeted therapy to the lungs. The invention could be used, for example, in such diverse applications as delivery of chemotherapeutic agents for treatment of lung cancer, or for systemic delivery of insulin in diabetics.

The references earlier mentioned are more fully identified hereafter, and are hereby incorporated by reference and relied upon:

- 10 Alton, E.W.F.W., Middleton, P.G., Caplen, N.J., Smith, S.N., Steel, D.M., Munkonge F.M., Jeffrey, P.K., Geddes, D.M., Hart, S.L., Williamson, R., Fasold, K.I., Miller, A.D., Dickson, P.,  
15 Stevenson, B.J., McLachlan, G., Dorin, J.R., and Porteous, D. (1993) *Nature Genetics* 5:135-142.
- Atkins, P.J., Barker, N.P. and Mathisen, D. (1992) In: *Pharmaceutical Inhalation Aerosol Technology* (Swarbrick, J., ed.) pp. 155-185, Marcel Decker, New York.
- 20 Bennett, C.F., Chiang, M.Y., Chan, H., Shoemaker, J.E., and Mirabelli, C.K. (1992) *Molec. Pharmacol.* 41: 1023-1033.
- Brigham, K.L., Meyrick, B., Christman, B., Magnuson, M., King, G., and Berry, L.C., Jr.  
25 (1989) *Am. J. Med. Sci.* 298:278-281.
- Byron, P.R. (1990) In: *Respiratory Drug Delivery* (Byron, P.R., ed.), pp. 167-206, CRC Press, Inc., Boca Raton, Florida.
- 30 Caplen, N.J., Alton, E.W.F.W., Middleton, P.G., Dorin, J.R., Stevenson, B.J., Gao, X., Durham, S.R., Jeffery, P.K., Hodson, M.F., Coutelle, C., Huang, L., Porteous, D.J., Williamson, R., and Geddes, D.M. (1995) *Nature Medicine* 1:39-46.
- 35 Caplen, N.J., Gao, X., Hayes, P., Elaswarapu, R., Fisher, G., Kinrade, E., Chakera, A., Schorr, J., Hughes, B., Dorin, J.R., Porteous, D.J., Alton, E.W.F.W., Geddes, D.M., Coutelle, C., Williamson,

- R., Huang, L., and Gilchrist, C. (1994) *Gene Therapy* 1:139-147.
- 5 Debs, R.J., Freedman, L.P., Edmunds, S., Gaensler, K.L., Duzgunes, N. and Yamamoto, K.R. (1990) *J. Biol. Chem.* 265:10189-10192.
- De Wet, J.R., Wood, K.V., DeLuca, M., Helinski, D.R., and Subramani, S. (1987) *Mol. Cell. Biol.* 7:725-737.
- 10 Farhood, H., Bottega, R., Epand, R.M., and Huang, L. (1992) *Biochim. Biophys. Acta* 1111:239-246.
- Farhood, H., Serbina, N., and Huang, L. (1995) *Biochim. Biophys. Acta*, In Press.
- Felgner, P.L., and Ringold, G.M. (1989) *Nature* 337:387-388.
- 15 Felgner, P.L., Gadek, T.R., Holm, M., Roman, R., Chan, H.W., Wenz, M., Northrop, J.P., Ringold, G.M., and Danielsen, M. (1987) *Proc. Natl. Acad. Sci. USA* 84:7413-7417
- 20 Gao, X., and Huang, L. (1991) *Biochem. Biophys. Res. Comm.* 179:280-285.
- Gorman, C.M., Moffat, L.F., and Howard, B.H. (1982) *Mol. Cell. Biol.* 2:1044-1051.
- Hickey, A.J. (1992) In: *Pharmaceutical Inhalation Aerosol Technology* (Swarbrick, J., ed.) pp. 167-206, Marcel Dekker, New York.
- 25 Hyde, S., Gill, D., Higgins, C.F., Trezise, A.E., MacVinish, L.J., Cuthbert, A.W., Ratcliff, R., Evans, M.J., and Colledge, W.H. (1993) *Nature* 362:250-255.
- 30 Logan, J.J., Bebok, Z., Walker, L.C., Peng, S., Felgner, P.L., Siegal, G.P., Frizzell, R.A., Dong, J., Howard, M. Matalon, S. Lindsey, J.R., DuVall, M., and Sorscher, E.J. (1995) *Gene Therapy* 2:38-49.
- 35 Malone, R.W., Felgner, P.L., and Verma, I.M. (1989) *Proc. Natl. Acad. Sci. USA* 86:6077-6081.

Morgan, R.A., and Anderson, W.F. (1993) *Ann. Rev. Biochem.* 62:191-217.

Nabel, G.J., Chang, E.G., Nabel, E.G., Plautz, G., Fox, B.A., Huang, L., and Shu, S. (1992). *Hum. Gene Ther.* 3:399-410.

Sambrook, J., Fritsch, E.F., and Maniatis, T. In *Molecular Cloning: A Laboratory Manual*, Vol. 1., Cold Springs Harbor Laboratory Press, 1989.

Zhu, N., Ligitt, D., Liu, Y., and Debs, R. (1993). *Science* 261:209-211.

While the invention has been described in connection with what is presently considered to be the most practical and preferred embodiment, it is to be understood that the invention is not to be limited to the disclosed embodiment, but is intended to cover various modifications included within the spirit and scope of the appended claims.



WHAT IS CLAIMED IS:

1. A dry powder formulation comprising a DNA/lipid complex, RNA/lipid complex or protein/lipid complex suitable for administration  
5 to an individual.
2. The formulation of claim 1 where the lipid portion of said complexes comprises 3 $\beta$ [N-(N',N'-dimethylaminoethane)-carbamoyl] cholesterol.
- 10 3. The formulation of claim 1 where the DNA encodes CFTR protein.
4. The formulation of claim 1 where a suitable amount of a cryoprotectant has been added.
- 15 5. The formulation of claim 4 where the cryoprotectant is a sugar.
6. The formulation of claim 5 where the sugar is lactose.
7. A method of preparing DNA, RNA,  
20 oligonucleotide or protein in powder form suitable for therapeutic administration to an individual, said method comprising mixing said DNA, RNA, oligonucleotide or protein with liposomes in an aqueous solution so that DNA/liposome,  
25 RNA/liposome, oligonucleotide/liposome or protein/liposome complexes are formed, and freeze-drying the resulting complexes.

8. The method of claim 7 which further comprises adding a cryoprotective agent prior to freeze-drying.

5 9. A method of delivering DNA, RNA, oligonucleotide or protein to an individual, said method comprising administration of the formulation of claim 1 to the individual.

10 10. The method of claim 9 where the formulation is administered to the nasal passages, trachea or lungs by means of an aerosol.

11. The method of claim 9 where the formulation is reconstituted in an aqueous solution and administered by nebulization, injection or infusion.

15 12. The method of claim 9 where the formulation is administered topically.

20 13. In a method of conducting gene therapy on a mammal in need of such therapy, wherein a DNA or RNA formulation is administered to the mammal, the improvement comprising administering the DNA or RNA as a dry powder formulation which contains a lyophilized DNA/liposome or RNA/liposome complex.

25 14. The method of claim 13 wherein the dry powder formulation contains a cryoprotective agent.

25

15. The method of claim 13 wherein the formulation is administered as an aerosol.


16. The formulation of claim 1 wherein said DNA or RNA comprises an oligonucleotide.


5           17. The formulation of claim 16 where a suitable amount of a cryoprotectant has been added.

18. The formulation of claim 17 where the cryoprotectant is a sugar.

10           19. The formulation of claim 18 where the sugar is lactose.



2 / 3  
FIGURE 2



## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US96/02681**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) :A61K 48/00

US CL :514/44; 435/320.1

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols,

U.S. : 514/44; 435/320.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, MEDLINE, BIOSIS, EMBASE, CAPLUS, WPIDS

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US, A, 5,384,128 (MEEZAN ET AL) 24 January 1995, see entire document.	1-19
Y	Chemical and Pharmacology Bulliten, Volume 40, Number 01, issued January 1992, TANAKA ET AL., "Cryoprotective Mechanism of Saccharides on Freeze-Drying of Liposome", pages 1-5, see entire document.	1-19
Y	Biochemical and Biophysical Research Communications, Volume 179, Number 1, issued 30 August 1991, GAO ET AL., "A Novel Cationic Liposome Reagent For Efficient Transfection of Mammalian Cells", pages 280-285, see entire document.	1-19

☐ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be part of particular relevance	X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (so specified)	A	document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means		
*P* document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

19 MAY 1996

Date of mailing of the international search report

12 JUN 1996

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

ANDREW MILNE

Telephone No. (703) 308-0196